

Glutathione – The Body’s Master Detoxifier and Antioxidant

James Meschino DC, MS, ND

Glutathione is a tripeptide comprised of the amino acids L-cysteine, L-glutamic and glycine. At the cellular level glutathione functions as a water-soluble antioxidant and is directly involved in specific detoxification reactions that protect the body against various dangerous substances. (1)

Glutathione, the most abundant cellular thiol, provides the major antioxidant defense mechanism in all mammalian cells by neutralizing toxic peroxides (e.g. hydrogen peroxide formed from oxygen metabolism). Glutathione is known to be consumed more rapidly, and depleted, upon ingestion of certain medications and environmental toxins. It is also depleted in cases of chronic liver infection and HIV infection. Evidence suggests that depletion of liver glutathione stores below a critical threshold level is the final event that triggers the conversion of HIV to the more ominous signs, symptoms and complications of AIDS. (2-4)

Unfortunately, for those wishing to boost their glutathione levels, it should be noted that glutathione supplements are not well absorbed from the intestinal tract. (5) However, supplementation with some standard antioxidants, along with N-acetylcysteine, alpha-lipoic acid, silymarin flavonoid (from milk thistle) and L-glutamine, are proven methods to support and raise serum and tissue levels of glutathione on a daily basis.

Nutritional support to boost glutathione levels should be considered as a useful adjunct in the management of HIV infection, chronic liver conditions, alcohol or drug-induced liver damage, and to combat exposure to environmental toxins, pollutants, medications, and certain carcinogens, and to help combat the age-related decline in detoxification ability that occurs after age 50.

Antioxidant Function

Glutathione is a premiere cellular antioxidant. It exists in reduced (GSH) and oxidized (GSSG) states within our cells. In the reduced state, the thiol group of cysteine is able to quench free radicals by donating a reducing equivalent ($H^+ + e^-$). This includes quenching various reactive oxygen species (ROS) that are generated during the use of oxygen by our cells in aerobic metabolism. In donating an electron, glutathione itself becomes reactive, but readily reacts with another reactive glutathione to form glutathione disulfide (GSSG). This is possible due to the normally high concentration of glutathione in cells (up to 5 mM in the liver). GSH can be regenerated from GSSG by the enzyme glutathione reductase.

In healthy cells and tissue, more than 90% of the total glutathione pool is in the reduced form (GSH) and less than 10% exists in the disulfide form (GSSG). An increased GSSG-to-GSH ratio is highly indicative of high levels of oxidative stress (presence of free radicals). As such, the activity of glutathione reductase is used as an indicator for oxidative stress. Glutathione status can be assessed by lab tests using Ellman's reagent or roGFP (redox-sensitive Green Fluorescent Protein).

Glutathione is also a cofactor for the enzyme glutathione peroxidase, which is a critical cellular antioxidant. It also helps to maintain levels of ascorbate and tocopherol by acting as a reducing agent. Conversely, vitamin C and vitamin E supplementation can also help to spare glutathione depletion. ((1,2,3,4,6,7,8,9, 21)

Detoxification Function

Glutathione conjugation is the primary mechanism of eliminating electrophilic (free radicals) xenobiotics (some of which are carcinogens) in the liver. Glutathione is required for conjugation

reactions and reduction reactions, catalyzed by glutathione S-transferase enzymes in the cytosol, microsomes, and mitochondria. The family of glutathione S-transferase enzymes are responsible for the quenching and detoxification of many environmental substances, including free radicals, peroxidized lipids, and xenobiotics (wide variety of environmental toxins, drugs and antibiotics). In the modern world, the liver's detoxification capacity can be easily overwhelmed, leading to glutathione depletion and an accumulation of toxins and potentially dangerous by-products of drug and antibiotic metabolism, due to high level of exposure to these man-made compounds. (10,11,12,13,14)

A quick historical review highlights the fact that since the Second World War an estimated 85,000 synthetic chemicals have been registered in the US alone. Today's environment now exposes us to various pesticides, herbicides, plastics (e.g. PVC), xenoestrogens (e.g. bisphenol A, parabens) and pollutants in our food, water and air. The human body has had very little time to adapt to these new challenges, many of which can over-tax our detoxification system, suppress immune function and exert other undesirable health effects. As well, many individuals are prescribed several drugs at one time (polypharmacy), which further depletes glutathione stores.

Many experts believe that the new chemical environment in which we live is a major contributing factor to the high incidence of many degenerative diseases in the modern era. (15)

As an example, glutathione is required for the safe detoxification of acetaminophen (paracetamol). In this instance it participates in a non-enzymatic conjugation of *N*-acetyl-*p*-benzoquinone imine (NAPQI), the reactive cytochrome P450- metabolite formed by acetaminophen that becomes toxic when GSH is depleted by an overdose of acetaminophen or its chronic use.

If NAPQI is permitted to build up in liver cells it reacts with cellular proteins, killing the cells in the process. As such, chronic use of acetaminophen is known to cause liver damage. This is largely due to the depletion of glutathione. Other drugs have been shown to deplete glutathione as well, rendering liver cells susceptible to damage and death from various drug by-products. (16, 17)

As such, health practitioners should ensure that patients employ strategies to guard against drug-induced liver damage from medications, by teaching patients how to re-constitute their liver glutathione stores on a daily basis.

Glutathione is also a vital substance that converts fat-soluble toxins into water-soluble forms, so that they can be safely and efficiently eliminated by the body. This helps prevent the accumulation in the body of various fat-soluble toxins. Glutathione is also needed for the detoxification of methylglyoxal, a toxin produced as a by-product of metabolism. (1,8)

Immune Function

Glutathione is necessary for maintaining immune mediated T-cell activation and phagocytosis, in addition to cellular and antibody mediated cytotoxicity. It is also required to maintain a normal balance between the T-helper cell 1 (IL-2, IL-12, gamma-interferon) and the T-helper cell 2 (IL-6, IL-4, tumor necrosis factor-alpha, IL-10, IL-1) cytokine response profile.

Plasma glutathione levels in HIV-infected individuals, even in the asymptomatic state, have been found to be depressed as early as three weeks post-infection. Intracellular glutathione levels in both infected CD4 and CD8 lymphocyte subsets are also significantly depressed; levels from 62-69 percent of normal have been found in the CD4 and CD8 lymphocytes of HIV and AIDS patients.

Depletion of glutathione of 10-40 percent is capable of completely inhibiting T-cell activation in vitro. Research assessing ratios of reduced-to-oxidized glutathione in HIV-positive patients found significantly increased levels of oxidized glutathione, compared to HIV-negative controls.

These disturbances were greater in patients with more advanced disease. Low serum thiol levels (precursors to glutathione) have been shown in HIV-infected, injecting drug users (IDU) are associated with an increased risk of mortality: IDU with low serum thiol levels are 5.65 times more likely to experience an accelerated time-to-death.

Glutathione appears to also slow progression of HIV infection by blocking activation of tissue necrosis factor alpha and inhibiting the activity of reverse transcriptase (a major enzyme necessary for HIV replication) by 80-90 percent in cell cultures.

As such, many HIV-infected patients take supplements that are known to raise their cellular and immune levels of glutathione – a factor that is associated with retarding disease progression. (1, 9, 10, 11, 12, 13, 14, 18, 19, 20)

Restoring and Maintaining Glutathione Stores

As a first step to preserving glutathione levels, adults should consider taking a high potency multiple vitamin and mineral containing the following dosages:

Vitamin C – 1000 mg

Vitamin E Succinate – 400 IU

Selenium 100-200 mcg

Beta-carotene – 10,000 -20,000 IU

Vitamin A – 2500 IU (21)

In addition, there are several nutrients that are proven to restore and boost glutathione levels to a significant degree, even in HIV-infected patients and other challenging cases. These nutrients include N-acetylcysteine, alpha-lipoic acid, silymarin (from milk thistle) and L- glutamine. I generally recommend that patients take a combination supplement, which contains all of these factors at meaningful dosages, to support their glutathione concentrations.

N-acetylcysteine (NAC)

N-acetylcysteine has been used successfully to treat hepatic and renal failure caused by glutathione depletion secondary to acetaminophen overdose. It may also help in the treatment of certain pulmonary disease, due to its mucolytic properties. N-acetylcysteine can also act as a heavy metal chelating agent to rid the body of certain toxic metals, as well as for copper, zinc, and boron. It has also been shown to elevate glutathione levels in immune compromised states, when taken as a supplement. Many HIV patients have used N-acetylcysteine as part of adjunctive nutritional management. N-acetylcysteine has also been shown to slow the progression of chronic kidney disease and may be very beneficial to diabetic patients in this regard.

Patients with chronic liver infections (e.g. hepatitis C) have also realized improvement in their condition when NAC was used as part of adjunctive nutritional management. (16,17,18,19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 51)

Alpha-Lipoic Acid

Lipoic acid is a water-soluble and fat soluble antioxidant that is also shown to raise glutathione levels and regenerate vitamin C and vitamin E. Alpha-lipoic acid has been successfully raised glutathione levels in HIV patients and is shown to block other steps in the progression of this

disease. A clinical trial in 11 AIDS-diagnosed patients taking 450 mg lipoic acid daily for 14 days resulted in increases in plasma ascorbate, total glutathione, total plasma thiols, decreases in lipid peroxide levels and a significant elevation in CD4 and CD4/CD8 ratios.

Lipoic acid has been shown to help diabetic and prediabetic patients by increasing insulin sensitivity and successfully treating early stage diabetic neuropathy. Alpha-lipoic acid protects the central nervous system from free radical damage, which makes it an important nutritional consideration in cases of neuropathy, as well as MS, ALS, Parkinson's disease and possibly Alzheimer's disease. (31,32,33, 34, 35, 36, 37, 38)

Milk Thistle

Milk Thistle contains silymarin, which is a mixture of flavonolignans consisting chiefly of silibinin, silidianin, and silicristin. It is essentially the silymarin content of Milk Thistle extract, which has been shown to provide its medicinal effects, especially in regards to the treatment and prevention of various liver conditions. In 1986, the German Commission E approved an oral extract of Milk Thistle standardized to 70% crude silymarin content as a treatment for liver disease.

Silymarin increases liver glutathione content by over 35 percent in healthy human subjects and by over 50 percent in rats. As noted previously, glutathione is not only an antioxidant, but is required for phase I and phase II detoxification function within the liver, facilitating the detoxification of many xenobiotics, drugs, and carcinogens. The ability of Milk Thistle to help restore liver glutathione levels is considered to be a primary means by which silymarin has been shown to be an effective treatment for various liver diseases.

Silibinin also stimulates RNA polymerase A (also known as polymerase I) and DNA synthesis, which in turn increase the synthesis of ribosome proteins and thus, stimulates cell development. This ultimately increases the regenerative capacity of liver cells and results in the production of new liver cells to replace the damaged old ones. As such, silymarin has been shown to aid in the repair of liver cells that have been damaged by various microorganisms, alcohol, and other damaging chemicals.

Silymarin has been shown to increase the concentrations of superoxide dismutase – a very powerful intracellular antioxidant, which quenches the superoxide anion (a very aggressive damaging and reactive, free radical oxygen species). (39, 40, 41, 42, 43, 44)

L Glutamine

L- Glutamine supplementation has been shown to enhance glutathione stores in conjunction with N-acetylcysteine. It has been shown to help protect the gastrointestinal tract from damage by certain chemotherapy drugs (i.e. fluorouracil) and also prevents diarrhea that these drugs are known to produce.

Glutamine supplementation has been shown to enhance immune system function and result in a lower level of infection and a shorter stay in hospital following surgery, radiation treatment, bone marrow transplantation, and in patients suffering from injury, compared with patients receiving Glutamine-free parenteral nutrition.

L-glutamine is found in the body in higher quantities and concentrations than any other free amino acid. Numerous studies show the benefits of L- glutamine in strengthening the body's immune system when individuals are under stress, including reducing the number of upper respiratory tract infections in athletes who are undergoing high intensity training.(45,46,47,48,49,50)

Summary

The many different environmental toxins, medications and other xenobiotics that are commonplace in our world today have been shown to more rapidly use up the body's pool of glutathione as per glutathione's involvement in quenching electrophiles (free radicals), neutralizing peroxides, and detoxifying numerous chemicals via phase I and phase II (conjugation reactions) pathways. The evidence is strong and consistent that glutathione depletion renders liver cells susceptible to the damaging effects of various drugs and other toxins, contributing to advancing liver disease.

Striving to ensure more optimal glutathione nutritional status appears to be a prudent strategy to help the body defend itself against liver damage and reduce concentrations of circulating carcinogens in the system. Boosting glutathione levels also helps to support immune function and provides important antioxidant protection to all body cells. Maintaining glutathione status is also important in the nutritional management of chronic liver infections and diseases and HIV infection.

The key nutrients that support glutathione status (N-acetylcysteine, alpha-lipoic acid, silymarin and L-glutamine) also provide additional health benefits such as protecting against nerve cell damage, treating neuropathies, repairing damaged liver cells, strengthening immune function, improving glucose tolerance, chelating heavy metals, supporting digestive health and slowing the progression of chronic kidney disease.

In my view, for general day-to-day support I suggest taking a combination capsule containing:

- N-Acetyl Cysteine – 350 mg
- Alpha – Lipoic acid – 75 mg
- L- Glutamine – 300 mg
- Milk Thistle (standardized to 80% silymarin content) – 100 mg

In regards to recommending this glutathione-boosting formulation to patients, practitioners may wish to consider the following evidence-based guidelines:

Clinical Applications and Dosage Considerations:

- General Health Support – 1 capsule per day
- Liver Ailments – 2-4 capsules per day
- Chronic Kidney Disease – 2-4 capsules per day
- Regular Acetaminophen or Alcohol Use – 1-2 capsules per day
- Patients taking two or more prescription drugs daily (polypharmacy) – 2 capsules per day
- HIV Infection – 3-4 capsules per day
- Bronchitis (to break up mucus) – 2 capsules per day
- Diabetes – (to improve glucose tolerance and antioxidant function) - 2 capsules per day
- Amyotrophic Lateral Sclerosis (ALS), Multiple Sclerosis, Parkinson's Disease - 4 capsules per day
- To Help Reverse Build Up Of Mercury, Cadmium, Arsenic and other heavy metals – 1-4 capsules per day (based on severity)
- Cataracts – 2 capsules per day
- Polycystic Ovarian Disease – 2 capsules per day
- Adjunctive Nutritional Cancer Management – 3-4 capsules per day

For more information on this or other related topics, please visit:

<http://www.meschinohealth.com>

References

1. Murray RK. Metabolism of xenobiotics. In: Murray RK, Granner DK, Mayes PA, et al, eds. Harper's Biochemistry. 24th ed. Stamford, CT: Appleton and Lange; 1996.
2. Carr A, Cooper DA, Penny R. Allergic manifestations of human immunodeficiency virus (HIV) infection. *J Clin Immunol* 1991;11:55-64.
3. Shriner K, Goetz MB. Severe hepatotoxicity in a patient receiving both acetaminophen and zidovudine. *Am J Med* 1992;93:94-96.
4. Schreck R, Rieber P, Baeuerle P. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *EMBO J* 1991;10:2247-2258.
4. Israel N, Gougerot-Pocidal MA, Aillet F, Virelizier JL. Redox status of cells influences constitutive or induced NF-kB translocation and HIV long terminal repeat activity in human T and monocytic cell lines. *J Immunol* 1992;149:3386-3393.
5. Witschi A, Reddy S, Stofer B, Lauterburg BH. The systemic availability of oral glutathione. *Eur J Clin Pharmacol* 1992;43:667-669.
6. Meister A, Anderson ME. Glutathione. *Annu Rev Biochem* 1983;52:711-760.
7. Meister A. Glutathione-ascorbic acid antioxidant system in animals. *J Biol Chem* 1994;269:9397-9400
8. Hamilos DL, Wedner HJ. The role of glutathione in lymphocyte activation. I. Comparison of inhibitory effects of buthionine sulfoximine and 2-cyclohexene-1-one by nuclear size transformation. *J Immunol* 1985;135:2740-2747.
9. Muller F, Aukrust P, Svardal AM, et al. The thiols glutathione, cysteine and homocysteine in immunodeficiency virus infection. In: Watson RR, ed. *Nutrition and Foods in AIDS*. Boca Raton, FL: CRC Press; 1998:35-39.
10. Marmor M, Alcabes P, Titus S, et al. Low serum thiol levels predict shorter times-to-death among HIV-infected injecting drug users. *AIDS* 1997; 11:1389-1393.
11. Buhl R, Jaffe HA, Holroyd KJ, et al. Systemic glutathione deficiency in symptom free HIV-seropositive individuals. *Lancet* 1989;2:1294-1298.
12. Staal FJ, Roederer M, Herzenberg LA, Herzenberg LA. Intracellular thiols regulate activation of nuclear factor kB and transcription of human immunodeficiency virus. *Proc Natl Acad Sci USA* 1990;87:9943-9947.
13. Aukrust P, Muller F. Glutathione redox disturbances in human immunodeficiency virus infection: immunologic and therapeutic consequences. *Nutrition* 1999; 15:165-167.
14. Buttke TM, Sandstrom PA. Oxidative stress as a mediator of apoptosis. *Immunol Today* 1994;15:7-10.

15. The Antioxidant Prescription. Bryce Wylde. Random House. 2008: 35
16. Smilkstein MJ, Knapp GL, Kulig KW, Rumack BH. Efficacy of oral N-acetylcysteine in the treatment of acetaminophen overdose. Analysis of the national multicenter study, (1976-1985). *N Engl J Med* 1988;319:1557-1562.
17. Azarnooof D, Arnold G J, Anderson J, et al. Acetylcysteine. In: *Mosby's Gen Rx*. St. Louis, MO: Mosby-Year Book Inc.; 1998.
18. Pirmohamed M, Williams D, Tingle MD, et al. Intracellular glutathione in the peripheral blood cells of HIV-infected patients: failure to show a deficiency. *AIDS* 1996;10:501-507.
19. de Quay B, Malinverni R, Lauterburg BH. Glutathione depletion in HIV-infected patients: role of cysteine deficiency and effect of oral N-acetylcysteine. *AIDS* 1992;6:815-819.
20. Kalebic T, Kinter A, Poli G, et al. Suppression of human immunodeficiency virus expression in chronically infected monocytic cells by glutathione, glutathione ester, and N-acetylcysteine. *Proc Natl Acad Sci USA* 1991;88:986-990.
21. *Encyclopedia of Natural Medicine* (revised 2nd edition) Murray M.: 204-205 (Prima Publishing 1998).
22. Smilkstein MJ, Knapp GL, Kulig KW, Rumack BH. Efficacy of oral N-acetylcysteine in the treatment of acetaminophen overdose. Analysis of the national multicenter study, (1976-1985). *N Engl J Med* 1988;319:1557-1562.
23. Azarnooof D, Arnold G J, Anderson J, et al. Acetylcysteine. In: *Mosby's Gen Rx*. St. Louis, MO: Mosby-Year Book Inc.; 1998.
24. Behr J, Maier K, Degenkolb B, et al. Antioxidative and clinical effects of high-dose N-acetylcysteine in fibrosing alveolitis. Adjunctive therapy to maintenance immunosuppression. *Am J Respir Crit Care Med* 1997;156:1897-1901.
25. Kelly G. Clinical applications of N-acetylcysteine. *Altern Med Rev* 1998;3:114-127.
26. De Flora S, Bennicelli C, Camoirano A, et al. In vivo effects of N-acetylcysteine on glutathione metabolism and on the biotransformation of carcinogenic and/or mutagenic compounds. *Carcinogenesis* 1985;6:1735-1745.
27. Hoffer E, Baum Y, Tabak A, Taitelman U. N-acetylcysteine increases the glutathione content and protects rat alveolar type II cells against paraquat-induced cytotoxicity. *Toxicol Lett* 1996;84:7-12.
28. Corcoran GB, Wong BK. Role of glutathione in prevention of acetaminophen-induced hepatotoxicity by N-acetyl-L-cysteine in vivo: studies with N-acetyl-D-cysteine in mice. *J Pharmacol Exp Ther* 1986;238:54-61.
29. Bernard GR, Wheeler AP, Arons MM, et al. A trial of antioxidants N-acetylcysteine and procysteine in ARDS. The Antioxidant in ARDS Study Group. *Chest* 1997;112:164-172.
30. Kinscherf R, Fischbach T, Mihm S, et al. Effect of glutathione depletion and oral N-acetylcysteine treatment on CD4 and CD8 cells. *FASEB J* 1994;8:448-451.

31. Packer L, Witt EH, Tritschler HJ. alpha-Lipoic acid as a biological antioxidant. *Free Radic Biol Med* 1995;19:227-250.
32. Packer L, Tritschler HJ, Wessel K. Neuroprotection by the metabolic antioxidant alpha-lipoic acid. *Free Radic Biol Med* 1997;22:359-378.
33. Han D, Tritschler HJ, Packer L. Alpha-lipoic acid increases intracellular glutathione in a human T-lymphocyte Jurkat cell line. *Biochem Biophys Res Commun* 1995;207:258-264.
34. Packer L, Suzuki YJ. Vitamin E and alpha-lipoate: role in antioxidant recycling and activation of the NF-kappa B transcription factor. *Mol Aspects Med* 1993;14:229-239.
35. Baur A, Harrer T, Peukert M, et al. Alpha-lipoic acid is an effective inhibitor of human immuno-deficiency virus (HIV- 1) replication. *Klin Wochenschr* 1991;69:722-724.
36. Fuchs J, Schofer H, Milbradt R, et al. Studies on lipoate effects on blood redox state in human immunodeficiency virus infected patients. *Arzneimittelforschung* 1993;43:1359-1362.
37. Kieburtz K, Schifitto G, McDermott M, et al. A randomized, double-blind, placebo-controlled trial of deprenyl and thioctic acid in human immunodeficiency virus-associated cognitive impairment. Dana Consortium on the Therapy of HIV Dementia and Related Cognitive Disorders. *Neurology* 1998;50:645-651.
38. Jayanthi S, Varalakshmi P. Tissue lipids in experimental calcium oxalate lithiasis and the effect of DL alpha-lipoic acid. *Biochem Int* 1992;26:913-921.
39. Hikino H, Kiso Y, Wagner H, Fiebig. Antihepatotoxic actions of flavanolignans from silybum marianum fruits. *Planta Medica* 1984;50:248-50
40. Muzes G, Deak G, Land I et al. Effect of the bioflavonoid silymarin on the in vitro activity and expression of superoxide dismutase (SOD) enzyme. *Acta Physiologica Hungarica* 1991;78(1):3-9
41. The Botanical Pharmacy. Quarry Health Books 2000. Boon H and Smith M:250
42. Natural Products Encyclopedia. www.consumerslab.com: Milk Thistle
43. Dietary Supplement Information Bureau. www.content.intramedicine.com: Thistle
44. Salmi HA, Sarna S. Effect of silymarin on chemical, functional and morphological alterations of the liver. A double-blind controlled study. *Scand J Gastroenterol* 1982;17:517-21
45. Miller AL. Therapeutic considerations of L-glutamine: a review of the literature. *Altern Med Rev* 1999;4:239-48.
46. Patrick L. Nutrients and HIV: part three N-acetylcysteine, alpha-lipoic acid, L-Lglutamine, and L-carnitine. *Altern Med Rev* 2000;4:290-305.
47. Daniele B, Perrone F, Gallo C, Pignata S, De Martino S, De Vivo R, et al. Oral glutamine in the prevention of fluorouracil induced intestinal toxicity: a double blind, placebo controlled, randomised trial. *Gut* 2001;48:28-33.

48. Griffiths RP. Outcome of critically ill patients after supplementation with glutamine. *Nutrition* 1997;13:752-4.

49. Chang WK, Yang KD, Shaio MF. Effect of glutamine on TH1 and TH2 cytokine responses of human peripheral blood mononuclear cells. *Clin Immunol* 1999;3:294-301.

50. Calder PC, Yaqoob P. Glutamine and the immune system. *Amino Acids* 1999;17;3:227-41.

51. Jo SH, Koo BK, Park JS, Kang HJ, Kim YJ, et al. N-acetylcysteine versus Ascorbic acid for preventing contrast-Induced nephropathy in patients with renal insufficiency undergoing coronary angiography NASPI study-a prospective randomized controlled trial. *Am Heart J.* 2009 ; 157(3):576-583

Please Note: Above Reference links were accessible when the article was published. However, respective third-party sites may change the structure and content of their websites at any time, we are unable to guarantee that our links will always be up to date. We apologize for the inconvenience.